

# RANDOM MATRIX THEORY AND DISCRETE MOMENTS OF THE RIEMANN ZETA FUNCTION

C.P. HUGHES

ABSTRACT. We calculate the discrete moments of the characteristic polynomial of a random unitary matrix, evaluated a small distance away from an eigenangle. Such results allow us to make conjectures about similar moments for the Riemann zeta function, and provide a uniform approach to understanding moments of the zeta function and its derivative.

## 1. INTRODUCTION

Let  $\zeta(s)$  be the Riemann zeta function, and denote its non-trivial zeros by  $\frac{1}{2} + i\gamma_n$ , with  $0 < \gamma_1 \leq \gamma_2 \leq \dots$ . (For simplicity, we will assume the Riemann Hypothesis, which says that  $\gamma_n \in \mathbb{R}$ ). It is known (see, for example, Titchmarsh's book [1] for details) that if  $N(T)$  is the number of zeros with  $0 < \gamma_n \leq T$  then

$$N(T) = \frac{T}{2\pi} \log \frac{T}{2\pi e} + \mathcal{O}(\log T).$$

Moments of the zeta function,

$$I_k(T) := \frac{1}{T} \int_0^T |\zeta(\tfrac{1}{2} + it)|^{2k} dt,$$

have long been of interest to number theorists, with it being widely believed that

$$\frac{1}{T} \int_0^T |\zeta(\tfrac{1}{2} + it)|^{2k} dt \sim f_k a(k) (\log T)^{k^2}$$

with

$$(1.1) \quad a(k) = \prod_{p \text{ prime}} \left(1 - \frac{1}{p}\right)^{k^2} \sum_{m=0}^{\infty} \left(\frac{\Gamma(m+k)}{m! \Gamma(k)}\right)^2 p^{-m},$$

and  $f_k$  being an integer (when  $k$  is integer) divided by  $(k^2)!$ , whose value was unknown apart from in a few cases. The known values of  $f_k$  are  $f_0 = 1$  (trivial),  $f_1 = 1$  (Hardy and Littlewood [2]), and  $f_2 = \frac{1}{12}$  (Ingham [3]). The value  $f_3 = \frac{42}{9!}$  has been conjectured by Conrey and Ghosh [4] and  $f_4 = \frac{24024}{16!}$  is a conjecture of Conrey and Gonek [5]. We should mention that  $a(k)$  given in (1.1) can be calculated for certain  $k$ :  $a(0) = a(1) = 1$  and  $a(-1) = a(2) = \frac{6}{\pi^2}$ .

In [6] Keating and Snaith argued that one can create a probabilistic model for the zeta function around height  $T$  using the characteristic polynomial of an  $N \times N$

unitary matrix chosen according to Haar measure, when

$$N = \log \frac{T}{2\pi}.$$

Setting

$$\begin{aligned} Z_U(\theta) &:= \det(I - e^{-i\theta}U) \\ &= \prod_{n=1}^N \left(1 - e^{i(\theta_n - \theta)}\right), \end{aligned}$$

and defining  $M_N(2k) := \mathbb{E}_N \{|Z_U(0)|^{2k}\}$  where  $\mathbb{E}_N$  denotes expectation with respect to Haar measure, they found that

$$\begin{aligned} (1.2) \quad M_N(2k) &= \prod_{j=1}^N \frac{\Gamma(j)\Gamma(j+2k)}{\Gamma(j+k)^2} \\ &\sim \frac{G^2(k+1)}{G(2k+1)} N^{k^2} \end{aligned}$$

as  $N \rightarrow \infty$  for fixed  $k$  subject to  $\Re(k) > -1/2$ , where  $G(\cdot)$  is the Barnes  $G$ -function.

By comparing with the known (and previously conjectured) values of  $f_k$ , they were led to conjecture that  $f_k = \frac{G^2(k+1)}{G(2k+1)}$ .

**Conjecture 1. (Keating and Snaith).** *For fixed  $k > -1/2$ , as  $T \rightarrow \infty$ ,*

$$\frac{1}{T} \int_0^T |\zeta(\tfrac{1}{2} + it)|^{2k} dt \sim \frac{G^2(k+1)}{G(2k+1)} a(k) \left(\log \frac{T}{2\pi}\right)^{k^2}$$

where  $G(\cdot)$  is the Barnes  $G$ -function and  $a(k)$  is given in (1.1)

Following this, Hughes, Keating and O'Connell [7] used the characteristic polynomial to model the discrete moments of the derivative of the zeta function,

$$J_k(T) := \frac{1}{N(T)} \sum_{0 < \gamma_n \leq T} |\zeta'(\tfrac{1}{2} + i\gamma_n)|^{2k}.$$

They calculated that for fixed  $k$  subject to  $\Re(k) > -3/2$ ,

$$\mathbb{E} \left\{ \frac{1}{N} \sum_{n=1}^N |Z'_U(\theta_n)|^{2k} \right\} \sim \frac{G^2(k+2)}{G(2k+3)} N^{k(k+2)}$$

as  $N \rightarrow \infty$ , and they used this to conjecture the asymptotic form  $J_k(T)$  should take for large  $T$ .

**Conjecture 2. (Hughes, Keating and O'Connell).** *If all the zeros of the zeta function are all simple, then for fixed  $k > -3/2$ , as  $T \rightarrow \infty$ ,*

$$\frac{1}{N(T)} \sum_{0 < \gamma_n \leq T} |\zeta'(\tfrac{1}{2} + i\gamma_n)|^{2k} \sim \frac{G^2(k+2)}{G(2k+3)} a(k) \left(\log \frac{T}{2\pi}\right)^{k(k+2)}$$

where  $a(k)$  is given by (1.1), and  $G(\cdot)$  is the Barnes  $G$ -function.

Again this conjecture is found to agree with all previously known results; when  $k = -1$  (a conjecture of Gonek [8]), when  $k = 0$  (trivial), and when  $k = 1$  (a theorem of Gonek [9] under RH). Also, extending a theorem due to Conrey, Ghosh and Gonek [10] (recorded in this paper as theorem 6) beyond its range of (proven) applicability, conjecture 7.1 states that  $J_2(T) \sim \frac{1}{1440\pi^2} \left(\log \frac{T}{2\pi}\right)^8$ , which agrees perfectly with conjecture 2.

It is striking that conjecture 1 and conjecture 2 have very similar form. The purpose of this paper is to unify them as special cases of one result.

In the next section we will show

$$\mathbb{E}_N \left\{ \left| Z_U \left( \theta_1 + \frac{y}{N} \right) \right|^{2k} \right\} \sim \frac{G^2(k+1)}{G(2k+1)} F_k(y) N^{k^2}$$

where  $F_k(y)$  is a certain function, independent of  $N$ , given in theorem 3.

We will then use this to conjecture that for  $L = \frac{1}{2\pi} \log \frac{T}{2\pi}$ ,

$$\frac{1}{N(T)} \sum_{0 < \gamma_n \leq T} \left| \zeta \left( \frac{1}{2} + i(\gamma_n + \alpha/L) \right) \right|^{2k} \sim \frac{G^2(k+1)}{G(2k+1)} a(k) F_k(2\pi\alpha) \left( \log \frac{T}{2\pi} \right)^{k^2}$$

and will show, in section 3, that this conjecture contains conjectures 1 and 2 as special cases ( $\alpha \rightarrow \infty$  and  $\alpha \rightarrow 0$  respectively). The conjecture is found to agree with a known result of Gonek [9], and the extension of the theorem due to Conrey, Ghosh and Gonek [10] cited above. These comparisons are discussed in section 3.1.

## 2. THE RANDOM MATRIX CALCULATION

**Theorem 3.** *For fixed  $k$  with  $\Re(k) > -1/2$ , and for  $x \leq AN$  with  $A < \pi$  constant,*

$$\mathbb{E}_N \left\{ \left| Z_U \left( \theta_1 + \frac{2x}{N} \right) \right|^{2k} \right\} = \frac{G^2(k+1)}{G(2k+1)} F_k(2x) N^{k^2} + \mathcal{O}(N^{k^2-1})$$

where

$$(2.1) \quad F_k(2x) = x^2 j_k(x)^2 + x^2 j_{k-1}(x)^2 - 2kx j_k(x) j_{k-1}(x)$$

where  $j_n(x)$  are the spherical Bessel functions of the first kind.

**Proof.** First note that

$$|Z_U(\theta_N + \beta)|^{2k} = |1 - e^{-i\beta}|^{2k} \prod_{n=1}^{N-1} |1 - e^{i(\theta_n - \theta_N - \beta)}|^{2k}$$

The average of this over all  $N \times N$  unitary matrices with Haar measure can be written ([11, 12]) as an  $N$ -fold integral

$$\begin{aligned} \mathbb{E}_N \left\{ \prod_{n=1}^{N-1} |1 - e^{i(\theta_n - \theta_N - \beta)}|^{2k} \right\} = \\ \frac{1}{N!(2\pi)^N} \int_{-\pi}^{\pi} \cdots \int_{-\pi}^{\pi} \prod_{1 \leq i < j \leq N} |e^{i\theta_i} - e^{i\theta_j}|^2 \prod_{n=1}^{N-1} |1 - e^{i(\theta_n - \theta_N - \beta)}|^{2k} d\theta_1 \cdots d\theta_N \end{aligned}$$

Putting all the  $j = N$  terms from the first product into the second, we see that

$$(2.2) \quad \mathbb{E}_N \left\{ |Z_U(\theta_N + \beta)|^{2k} \right\} = \frac{1}{N} |2 \sin(\tfrac{1}{2}\beta)|^{2k} \mathbb{E}_{N-1} \left\{ |Z_{\tilde{U}}(0)|^2 |Z_{\tilde{U}}(\beta)|^{2k} \right\},$$

where  $Z_{\tilde{U}}$  is the characteristic polynomial of an  $(N-1) \times (N-1)$  unitary matrix.

By rotation invariance of Haar measure,

$$\mathbb{E}_{N-1} \left\{ |Z_{\tilde{U}}(0)|^2 |Z_{\tilde{U}}(\beta)|^{2k} \right\} = \mathbb{E}_{N-1} \left\{ |Z_{\tilde{U}}(0)|^{2k} |Z_{\tilde{U}}(\beta)|^2 \right\}$$

This is calculated in Theorem 4 below, where it is shown that for  $\Re(k) > -1/2$ ,

$$\begin{aligned} \mathbb{E}_N \left\{ |Z_U(0)|^{2k} |Z_U(y/N)|^2 \right\} \\ \sim \frac{G^2(k+1)}{G(2k+1)} \sum_{p=0}^{\infty} \frac{k(k-1+p)!(k+p)!}{p!(2k+p)!(2k+1+2p)!} (-1)^p y^{2p} N^{(k+1)^2} \end{aligned}$$

and substituting this into (2.2) (where we put  $\beta = y/N$ ) we see that

$$\mathbb{E}_N \left\{ \left| Z_U \left( \theta_N + \frac{y}{N} \right) \right|^{2k} \right\} = \frac{G^2(k+1)}{G(2k+1)} F_k(y) N^{k^2} \left( 1 + \mathcal{O} \left( \frac{1}{N} \right) \right)$$

where

$$(2.3) \quad F_k(y) = k \sum_{p=0}^{\infty} \frac{(k-1+p)!(k+p)!}{p!(2k+p)!(2k+1+2p)!} (-1)^p y^{2k+2p}$$

The spherical Bessel functions of the first kind are defined as

$$\begin{aligned} j_n(z) &= \sqrt{\frac{\pi}{2z}} J_{n+1/2}(z) \\ &= \sum_{m=0}^{\infty} \frac{(-1)^m (n+m)!}{m! (2n+2m+1)! 2^{2m}} (2z)^{n+2m} \end{aligned}$$

where  $J_\nu(z)$  is the  $\nu$ -th order Bessel function of the first kind. Hence,

$$F_k(2x) = x^2 j_k(x)^2 + x^2 j_{k-1}(x)^2 - 2kx j_k(x) j_{k-1}(x)$$

which can be seen by comparing the Taylor expansions.

The above large- $N$  asymptotics are for  $x = o(N)$ . This can be extended to  $|x| \leq AN$  for  $A < \pi$  an arbitrary constant as follows: Let  $\beta$  be a fixed constant subject to  $0 < \beta < 2\pi$ . By (2.2) and the results of Basor [13],

$$\begin{aligned} \mathbb{E}_N \left\{ |Z_U(\theta_1 + \beta)|^{2k} \right\} &= \frac{1}{N} |2 \sin(\tfrac{1}{2}\beta)|^{2k} \mathbb{E}_{N-1} \left\{ |Z_{\tilde{U}}(0)|^{2k} |Z_{\tilde{U}}(\beta)|^2 \right\} \\ &\sim \frac{G^2(k+1)}{G(2k+1)} N^{k^2}. \end{aligned}$$

If one lets  $x = N\beta/2$  then the large- $x$  asymptotics of Bessel functions (see, for example, chapter 9 of [14]) implies that

$$F_k(N\beta) = 1 + \mathcal{O} \left( \frac{1}{N} \right).$$

and so theorem 3 gives the correct first order term as  $N \rightarrow \infty$ , for  $|x| \leq AN$  with  $A < \pi$  a constant.  $\square$

**Remark.** When  $n$  is an integer,

$$j_n(x) = (-1)^n x^n \left( \frac{1}{x} \frac{d}{dx} \right)^n \frac{\sin x}{x}$$

which leads to a neat evaluation of  $F_k(2x)$  for integer  $k$ , the first few being:

$$(2.4) \quad \begin{aligned} F_1(2x) &= \frac{x^2 - \sin^2(x)}{x^2} \\ F_2(2x) &= \frac{x^4 - 3x^2 + 3x \sin(2x) + (2x^2 - 3) \sin^2(x)}{x^4} \\ F_3(2x) &= \frac{x^6 - 3x^4 - 45x^2 + (-12x^3 + 45x) \sin(2x) + (-3x^4 + 72x^2 - 45) \sin^2(x)}{x^6} \end{aligned}$$

**Theorem 4.**

$$\begin{aligned} \mathbb{E}_N \{ |Z(0)|^{2k} |Z(\beta)|^2 \} &= M_N(2k) N! (N+2k)! \times \\ &\times \sum_{n=0}^N \frac{(2 \sin(\beta/2))^{2n}}{n! (2k+n)!} \sum_{m=0}^{N-n} \frac{(N+k-m)! (k+m+n)!}{(N-n-m)! m!} e^{i\beta(2m-N+n)} \end{aligned}$$

where  $M_N(2k)$  is given in (1.2). If  $\frac{y}{N} \rightarrow 0$  as  $N \rightarrow \infty$ , then for  $\Re(k) > -1/2$ ,

$$\begin{aligned} \mathbb{E}_N \left\{ |Z(0)|^{2k} \left| Z\left(\frac{y}{N}\right) \right|^2 \right\} &= \frac{G^2(k+1)}{G(2k+1)} \sum_{p=0}^{\infty} \frac{k(k-1+p)! (k+p)!}{p! (2k+p)! (2k+1+2p)!} (-1)^p y^{2p} \\ &\times N^{(k+1)^2} \left( 1 + \mathcal{O}\left(\frac{1}{N}\right) \right) \end{aligned}$$

**Proof.** If  $k$  is an integer, then

$$\begin{aligned} \mathbb{E}_N \left\{ \prod_{n=1}^N |1 - e^{i\theta_n}|^{2k} |1 - e^{i\theta_n} e^{-i\beta}|^2 \right\} \\ = \frac{1}{(2\pi)^N N!} \int \cdots \int_{-\pi}^{\pi} \prod_{1 \leq j < m \leq N} (e^{i\theta_j} - e^{i\theta_m})(e^{-i\theta_j} - e^{-i\theta_m}) \\ \times \prod_{n=1}^N (1 - e^{i\theta_n})^k (1 - e^{-i\theta_n})^k (1 - e^{i\theta_n} e^{-i\beta})(1 - e^{-i\theta_n} e^{i\beta}) d\theta_n \end{aligned}$$

which equals, after some simple manipulation of the terms

$$\begin{aligned} &\frac{(-1)^{N(N-1)/2+kN+N} e^{-iN\beta}}{(2\pi)^N N!} \times \\ &\int \cdots \int_{-\pi}^{\pi} \prod_{1 \leq j < m \leq N} (e^{i\theta_j} - e^{i\theta_m})^2 \prod_{n=1}^N (e^{i\theta_n})^{-N-k} (1 - e^{i\theta_n})^{2k} (e^{i\theta_n} - e^{i\beta})^2 d\theta_n \end{aligned}$$

Therefore,

$$\begin{aligned} \mathbb{E}_N \{ |Z_U(0)|^{2k} |Z_U(\beta)|^2 \} &= \frac{e^{-iN\beta}}{N!} (-1)^{N(N-1)/2+kN+N} \\ &\times \text{CT} \left\{ \prod_{1 \leq j < m \leq N} (t_j - t_m)^2 \prod_{n=1}^N \frac{1}{t_n^{N+k}} (1 - t_n)^{2k} (t_n - e^{i\beta})^2 \right\} \end{aligned}$$

where  $\text{CT}\{\cdot\}$  denotes the constant term in the Laurent expansion in the variables  $t_1, \dots, t_N$ . The constant term equals (by lemma 1 of [15])

$$\lim_{y \rightarrow 0} y^N \int \cdots \int_0^1 \prod_{1 \leq j < m \leq N} (t_j - t_m)^2 \prod_{n=1}^N t_n^{-N-k+y-1} (1 - t_n)^{2k} (t_n - e^{i\beta})^2 dt_n$$

Kaneko [16] has evaluated this integral (which is a generalization of Selberg's integral) as

$$\prod_{j=1}^N \frac{\Gamma(1+j)\Gamma(j+y-N-k)\Gamma(j+2k+1)}{\Gamma(j+y+k+1)} \sum_{m,n=0}^{\infty} \frac{(-N)_{m+n}(y+k+1)_{m+n}}{(y-N-k)_m(2k+1)_n} \frac{e^{2mi\beta}(1-e^{i\beta})^{2n}}{m!n!}$$

where  $(a)_n = a(a+1)\cdots(a+n-1) = \Gamma(a+n)/\Gamma(a)$ . Since we have assumed that  $k$  is an integer,

$$\lim_{y \rightarrow 0} y\Gamma(y+j-N-k) = \frac{(-1)^{N+k-j}}{\Gamma(N+k-j+1)}$$

and so we have

$$\begin{aligned} \mathbb{E}_N \{|Z_U(0)|^{2k}|Z_U(\beta)|^2\} &= \frac{e^{-iN\beta}}{N!} \prod_{j=1}^N \frac{\Gamma(1+j)\Gamma(j+2k+1)}{\Gamma(j+k+1)\Gamma(N+k-j+1)} \\ &\quad \times \sum_{m,n=0}^{\infty} \frac{(-N)_{m+n}(k+1)_{m+n}}{(-N-k)_m(2k+1)_n} \frac{e^{2mi\beta}(1-e^{i\beta})^{2n}}{m!n!} \end{aligned}$$

Expanding everything out in terms of the gamma function:

$$\begin{aligned} \mathbb{E}_N \{|Z_U(0)|^{2k}|Z_U(\beta)|^2\} &= M_N(2k) \frac{\Gamma(N+1+2k)N!}{\Gamma(N+1+k)^2} \\ &\quad \times \sum_{n=0}^N \frac{(2\sin(\beta/2))^{2n}}{n!\Gamma(2k+n+1)} \sum_{m=0}^{N-n} \frac{\Gamma(N+k+1-m)\Gamma(k+n+1+m)}{\Gamma(N+1-n-m)!} \frac{e^{i\beta(2m-N+n)}}{m!} \end{aligned}$$

where  $M_N(2k)$  is defined in (1.2). Observe that the inner summand is invariant as  $m \rightarrow N-n-m$ , and so the inner sum is in fact a sum of cosines (and thus the series expansion in  $\beta$  contains only even powers of  $\beta$ ). Furthermore observe that both sides of the equation are analytic functions of  $k$  (for  $\Re(k) \geq 0$ ), both sides can be easily bounded by  $\mathcal{O}(2^{2N\Re(k)})$  (for large  $k$ , with  $N$  and  $\beta$  fixed), and the two sides are equal at the positive integers. Thus by Carlson's theorem (see §17 of [12], for example), the restriction on  $k$  being an integer is no longer required, and the left-hand side equals the right-hand side for all complex  $k$ .

Now, using the fact that  $(-a)_n = (-1)^n \Gamma(a+1)/\Gamma(a+1-n)$ , we have

$$\begin{aligned} &\sum_{m=0}^{N-n} \frac{\Gamma(N+k+1-m)\Gamma(k+n+1+m)}{\Gamma(N+1-n-m)} \frac{e^{i\beta(2m-N+n)}}{m!} \\ &= \frac{\Gamma(N+k+1)\Gamma(k+n+1)}{\Gamma(N-n+1)} {}_2F_1 \left( \begin{matrix} -N+n, k+n+1 \\ -N-k \end{matrix}; e^{2i\beta} \right) e^{i(n-N)\beta} \end{aligned}$$

where

$${}_2F_1 \left( \begin{matrix} a, b \\ c \end{matrix}; z \right) = \sum_{m=0}^{\infty} \frac{(a)_m (b)_m}{(c)_m} \frac{z^m}{m!}$$

is the Gauss hypergeometric function. (If  $a, c$  are negative integers with  $-a < -c$  then it is a polynomial of degree  $-a$ ).

Applying the quadratic hypergeometric transformation 15.3.26 of [14] we get

$$(2.5) \quad {}_2F_1 \left( \begin{matrix} -N + n, k + n + 1 \\ -N - k \end{matrix}; e^{2i\beta} \right) e^{i(n-N)\beta} \\ = {}_2F_1 \left( \begin{matrix} -\frac{1}{2}N + \frac{1}{2}n, -\frac{1}{2}N + \frac{1}{2}n + \frac{1}{2} \\ -N - k \end{matrix}; \frac{1}{\cos^2(\beta)} \right) (2 \cos \beta)^{N-n}$$

For  $m$  a positive integer,

$${}_2F_1 \left( \begin{matrix} -m, b \\ c \end{matrix}; z \right) = \frac{\Gamma(1-c-m)\Gamma(1-c+b)}{\Gamma(1-c)\Gamma(1-c-m+b)} {}_2F_1 \left( \begin{matrix} -m, b \\ 1-c-m+b \end{matrix}; 1-z \right)$$

and so we see that the right-hand side of (2.5) equals

$$\frac{\Gamma(\frac{1}{2}N + \frac{1}{2}n + k + 1)\Gamma(\frac{1}{2}N + \frac{1}{2}n + k + \frac{3}{2})}{\Gamma(N + k + 1)\Gamma(k + n + \frac{3}{2})} (2 \cos \beta)^{N-n} \times \\ \times {}_2F_1 \left( \begin{matrix} -\frac{1}{2}N + \frac{1}{2}n, -\frac{1}{2}N + \frac{1}{2}n + \frac{1}{2} \\ n + k + \frac{3}{2} \end{matrix}; 1 - \frac{1}{\cos^2(\beta)} \right)$$

Therefore, we have proven

$$\mathbb{E}_N \{ |Z_U(0)|^{2k} |Z_U(\beta)|^2 \} = \\ \sum_{n=0}^N \sum_{m=0}^{\lfloor \frac{1}{2}(N-n) \rfloor} T(N, k, m, n) (-1)^m (2 \sin(\beta/2))^{2n} (\sin \beta)^{2m} (\cos \beta)^{N-n-2m}$$

where

$$T(N, k, m, n) = M_N(2k) \frac{N!(N+2k)! (\frac{1}{2}N + \frac{1}{2}n + k)! (\frac{1}{2}N + \frac{1}{2}n + k + \frac{1}{2})!}{(N-n)!(N+k)!^2} \\ \times \frac{2^{N-n} (-\frac{1}{2}N + \frac{1}{2}n)_m (-\frac{1}{2}N + \frac{1}{2}n + \frac{1}{2})_m (k+n)!}{m!n!(2k+n)!(n+k+\frac{1}{2}+m)!}$$

Observe that for fixed  $k, m, n$  with  $\Re(k) > -1/2$ ,

$$T(N, k, m, n) = \frac{G^2(k+1)}{G(2k+1)} \frac{(k+n)!(k+n+m)!}{m!n!(2k+n)!(2k+2n+2m+1)!} \\ \times N^{(k+1)^2+2n+2m} \left( 1 + \mathcal{O}\left(\frac{1}{N}\right) \right)$$

and so

$$\mathbb{E}_N \{ |Z(0)|^{2k} |Z(y/N)|^2 \} \sim \frac{G^2(k+1)}{G(2k+1)} N^{(k+1)^2} \\ \times \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} (-1)^m \frac{(k+n)!(k+n+m)!}{m!n!(2k+n)!(2k+2n+2m+1)!} y^{2m+2n}$$

in the sense that for each fixed integer  $h$ , the coefficient of  $y^h$  on the Taylor expansion of the left-hand side converges to that of the right-hand side as  $N \rightarrow \infty$ .

Finally we show that

$$\begin{aligned} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} (-1)^m \frac{(k+n)!(k+n+m)!}{m!n!(2k+n)!(2k+2n+2m+1)!} y^{2m+2n} \\ = \sum_{p=0}^{\infty} \frac{k(k-1+p)!(k+p)!}{p!(2k+p)!(2k+1+2p)!} (-1)^p y^{2p} \end{aligned}$$

This can be proven by comparing the coefficients of  $y^{2p}$ . That is, we wish to show for all integer  $p \geq 0$ ,

$$\sum_{n=0}^p \frac{(-1)^{p-n} (k+n)!(k+p)!}{(p-n)!n!(2k+n)!(2k+2p+1)!} = \frac{(-1)^p k(k-1+p)!(k+p)!}{p!(2k+p)!(2k+1+2p)!}$$

This is equivalent to showing

$$(2.6) \quad \sum_{n=0}^p \frac{(-1)^n (k+n)!p!(2k+p)!}{k(p-n)!n!(2k+n)!(k-1+p)!} = 1$$

for all integer  $p \geq 0$ , and this we shall do by creating the Wilf-Zeilberger pair [17]: Denote the summand in (2.6) by  $F(p, n)$ , and observe that

$$(2.7) \quad F(p+1, n) - F(p, n) = G(p, n+1) - G(p, n)$$

where

$$G(p, n) = \frac{(2k+n)n}{(n-p-1)(k+p)} F(p, n)$$

(the  $(2k+n)n/(n-p-1)(k+p)$  being calculated by Zeilberger's algorithm). Summing both sides of (2.7) over  $n$ , we see that the right-hand side telescopes to zero, which shows that the left-hand side of (2.6) must be a constant, independent of  $p$ . Putting  $p = 0$ , direct calculation shows that constant is 1.  $\square$

### 3. CONJECTURE ABOUT THE ZETA FUNCTION

**Conjecture 5.** *For fixed  $k$  subject to  $\Re(k) > -1/2$ ,*

$$\frac{1}{N(T)} \sum_{0 < \gamma_n \leq T} |\zeta(\tfrac{1}{2} + i(\gamma_n + \alpha/L))|^{2k} \sim \frac{G^2(k+1)}{G(2k+1)} a(k) F_k(2\pi\alpha) \left( \log \frac{T}{2\pi} \right)^{k^2}$$

as  $T \rightarrow \infty$ , uniformly in  $\alpha$  for  $|\alpha| \leq L$ , where  $L = \frac{1}{2\pi} \log \frac{T}{2\pi}$  is the density of zeros of height  $T$ .  $G(\cdot)$  is the Barnes  $G$ -function,  $a(k)$  is given by (1.1), and  $F_k(2\pi\alpha)$  is given in theorem 3.

If this conjecture is true, then we are able to prove conjecture 2 and a variant of the Keating-Snaith conjecture (conjecture 1).

**Corollary 5.1.** *If conjecture 5 is true, then*

$$\frac{1}{N(T)} \sum_{0 < \gamma_n \leq T} |\zeta'(\tfrac{1}{2} + i\gamma_n)|^{2k} \sim \frac{G^2(k+2)}{G(2k+3)} a(k) \left( \log \frac{T}{2\pi} \right)^{k(k+2)}$$



**Proof.** By the definition of differentiation,

$$\left| \zeta' \left( \frac{1}{2} + i\gamma_n \right) \right|^{2k} = L^{2k} \lim_{\alpha \rightarrow 0} \frac{\left| \zeta \left( \frac{1}{2} + i \left( \gamma_n + \frac{\alpha}{L} \right) \right) \right|^{2k}}{\alpha^{2k}}$$

From (2.3) we have

$$\lim_{\alpha \rightarrow 0} \frac{F(2\pi\alpha)}{\alpha^{2k}} = (2\pi)^{2k} \frac{k!k!}{(2k)!(2k+1)!}$$

so applying conjecture 5 and using uniformity to swap the  $\alpha \rightarrow 0$  and  $N \rightarrow \infty$  limits, we have

$$\begin{aligned} \frac{1}{N(T)} \sum_{0 < \gamma_n \leq T} \left| \zeta' \left( \frac{1}{2} + i\gamma_n \right) \right|^{2k} &\sim \frac{G^2(k+1)}{G(2k+1)} a(k) L^{2k} \frac{(2\pi)^{2k} k!k!}{(2k)!(2k+1)!} \left( \log \frac{T}{2\pi} \right)^{k^2} \\ &= \frac{G^2(k+2)}{G(2k+3)} a(k) \left( \log \frac{T}{2\pi} \right)^{k(k+2)} \end{aligned}$$

as required.  $\square$

**Corollary 5.2.** *From conjecture 5 it follows that for  $\beta > 0$  fixed*

$$(3.1) \quad \frac{1}{N(T)} \sum_{0 < \gamma_n \leq T} \left| \zeta \left( \frac{1}{2} + i(\gamma_n + \beta) \right) \right|^{2k} \sim \frac{G^2(k+1)}{G(2k+1)} a(k) \left( \log \frac{T}{2\pi} \right)^{k^2}$$

**Proof.** The asymptotics as  $z \rightarrow \infty$  of the spherical Bessel function (see chapters 9 and 10 of [14]) are

$$j_n(z) \sim \begin{cases} \frac{1}{z} (-1)^{(n+1)/2} \cos z & \text{if } n \text{ is odd} \\ \frac{1}{z} (-1)^{n/2} \sin z & \text{if } n \text{ is even} \end{cases}$$

and putting this into (2.1) we have

$$\lim_{\alpha \rightarrow \infty} F_k(2\pi\alpha) = 1$$

Setting  $\alpha = L\beta$  in conjecture 5 completes the proof.  $\square$

**Remark.** Note that corollary 5.2 can be thought of as a variant of conjecture 1. This is because one expects the mean of  $|\zeta(1/2 + it)|^{2k}$  to be independent of the average taken, and the Keating-Snaith conjecture is a result about the continuous mean, whereas corollary 5.2 is a result about a discrete mean. To see this, recall that the zeros get denser higher up the critical line, and so if  $\beta > 0$  is fixed and  $\gamma_n$  is random, one might expect  $\zeta \left( \frac{1}{2} + i(\gamma_n + \beta) \right)$  to be random (whereas, if  $\beta$  was small, it would be highly influenced by the fact that  $\zeta(1/2 + i\gamma_n) = 0$ ). The left-hand side of (3.1) averages this, and thus acts like a discrete mean of  $|\zeta(1/2 + it)|^{2k}$ .

**3.1. Comparison with the zeta function.** Gonek [9] showed that if the Riemann Hypothesis is true then

$$\frac{1}{N(T)} \sum_{0 < \gamma_n \leq T} \left| \zeta \left( \frac{1}{2} + i(\gamma_n + \alpha/L) \right) \right|^2 \sim \left( 1 - \left( \frac{\sin(\pi\alpha)}{\pi\alpha} \right)^2 \right) \log \frac{T}{2\pi}$$

uniformly in  $\alpha$  for  $|\alpha| \leq L/2$ , which is in perfect agreement with conjecture 5 when  $k = 1$ .

There is no proof of the conjecture for  $k = 2$  (unlike conjecture 1 which is proven for  $k = 1$  and 2). But there are theorems along the lines of conjecture 5 for  $k = 2$ :

**Theorem 6. (Conrey, Ghosh and Gonek [10].)** *Assume GRH and let  $A(s) = \sum_{n \leq x} n^{-s}$  where  $x = \left(\frac{T}{2\pi}\right)^\eta$  for some  $\eta \in (0, \frac{1}{2})$ . Then,*

$$\frac{1}{N(T)} \sum_{0 < \gamma_n \leq T} \left| \zeta A\left(\frac{1}{2} + i(\gamma_n + \alpha/L)\right) \right|^2 \sim \frac{6}{\pi^2} \sum_{j=0}^{\infty} \frac{(-1)^{j+1} (2\pi\alpha)^{2j+2}}{(2j+5)!} \times \\ \times \left( -\eta^2 + \frac{1}{3}(2j+5)\eta^3 - \frac{2j+5}{j+3}\eta^{2j+6} + \eta^{2j+7} + \eta^2(1-\eta)^{2j+5} \right) \left( \log \frac{T}{2\pi} \right)^4$$

uniformly for bounded  $\alpha$ .

(We have slightly changed notation from [10], to be consistent with our definition of  $L = \frac{1}{2\pi} \log \frac{T}{2\pi}$ ).

Putting  $\eta = 1$  in the above (which, as it stands, is not allowed under the conditions of the theorem) then  $A(\frac{1}{2} + it) = \zeta(\frac{1}{2} + it) + \mathcal{O}(t^{-1/2})$ , and we have

$$\frac{1}{N(T)} \sum_{0 < \gamma_n \leq T} \left| \zeta^2 \left( \frac{1}{2} + i(\gamma_n + \alpha/L) \right) \right|^2 \\ \sim \frac{4}{\pi^2} \sum_{j=1}^{\infty} \frac{(-1)^{j+1} (2\pi\alpha)^{2j+2}}{(2j+6)!} (2j^2 + 5j) \left( \log \frac{T}{2\pi} \right)^4$$

Note that

$$\frac{4}{\pi^2} \sum_{j=1}^{\infty} \frac{(-1)^{j+1} (2\pi\alpha)^{2j+2}}{(2j+6)!} (2j^2 + 5j) = \\ \frac{1}{12} a(2) \frac{(2\pi^2\alpha^2 - 3) \sin^2(\pi\alpha) + 3\pi\alpha \sin(2\pi\alpha) + (\pi\alpha)^4 - 3(\pi\alpha)^2}{(\pi\alpha)^4}$$

which is what is predicted in conjecture 5.

That is, from a purely number theoretical calculation involving no random matrix theory, we have

**Conjecture 7.** *Assuming that  $\eta = 1$  is permissible in theorem 6 then*

$$\frac{1}{N(T)} \sum_{0 < \gamma_n \leq T} \left| \zeta \left( \frac{1}{2} + i(\gamma_n + \alpha/L) \right) \right|^4 \sim \frac{1}{2\pi^2} F_2(2\pi\alpha) \left( \log \frac{T}{2\pi} \right)^4$$

where  $F_2(2x)$  is given in (2.4).

So, following the proof of corollary 5.1, we may deduce

**Corollary 7.1.** *If conjecture 7 is true then*

$$\frac{1}{N(T)} \sum_{0 < \gamma_n \leq T} \left| \zeta' \left( \frac{1}{2} + i\gamma_n \right) \right|^4 \sim \frac{1}{1440\pi^2} \left( \log \frac{T}{2\pi} \right)^8.$$

Note that this is the same answer that one gets from putting  $k = 2$  into conjecture 2.

## 4. ACKNOWLEDGMENTS

Early versions of the results described in this paper formed part of my PhD thesis [18], and I would like to thank my advisors, Jon Keating and Neil O’Connell, for their help and advice. Part of this research was carried out while I was supported by the EC TMR network “Mathematical aspects of Quantum Chaos”, EC-contract no HPRN-CT-2000-00103.

## REFERENCES

- [1] E.C. Titchmarsh, *The Theory of the Riemann Zeta-Function* 2nd ed., revised by D.R. Heath-Brown (Oxford Science Publications, 1986)
- [2] G.H. Hardy and J.E. Littlewood, “Contributions to the theory of the Riemann zeta-function and the theory of the distribution of primes”, *Acta Arith.* **41** (1918) 119–196
- [3] A.E. Ingham, “Mean-values theorems in the theory of the Riemann zeta-function”, *Proc. Lond. Math. Soc.* **27** (1926) 273–300
- [4] J.B. Conrey and A. Ghosh, “A conjecture for the sixth power moment of the Riemann zeta-function”, *Internat. Math. Res. Notices* **15** (1998) 775–780
- [5] J.B. Conrey and S.M. Gonek, “High moments of the Riemann zeta function”, *Duke Math. J.* **107** (2001) 577–604
- [6] J.P. Keating and N.C. Snaith, “Random matrix theory and  $\zeta(1/2 + it)$ ”, *Comm. Math. Phys.* **214** (2000) 57–89
- [7] C.P. Hughes, J.P. Keating and N. O’Connell, “Random matrix theory and the derivative of the Riemann zeta function”, *Proc. R. Soc. Lond. A* **456** (2000) 2611–2627
- [8] S.M. Gonek, “On negative moments of the Riemann zeta-function”, *Mathematika* **36** (1989) 71–88
- [9] S.M. Gonek, “Mean values of the Riemann zeta function and its derivatives”, *Invent. Math.* **75** (1984) 123–141
- [10] J.B. Conrey, A. Ghosh and S.M. Gonek, “Large gaps between zeros of the zeta function”, *Mathematika* **33** (1986) 212–238
- [11] H. Weyl, *Classical Groups*, (Princeton University Press, 1946)
- [12] M.L. Mehta, *Random Matrices* (Academic Press, 1991)
- [13] E.L. Basor, “Asymptotic formulas for Toeplitz determinants”, *Trans. Amer. Math. Soc.* **239** (1978) 33–65
- [14] M. Abramowitz and I.A. Stegun, *Handbook of Mathematical Functions* (Dover, ninth printing 1972)
- [15] P.J. Forrester, “Selberg correlation integrals and  $1/r^2$  quantum many-body system, *Nuclear Physics B* **388** (1992) 671–699
- [16] J. Kaneko, “Selberg integrals and hypergeometric functions associated with Jack polynomials”, *SIAM J. Math. Anal.* **24** (1993) 1086–1110
- [17] M. Petkovšek, H. Wilf and D. Zeilberger, *A=B* (A.K. Peters, 1996)
- [18] C.P. Hughes, *On the characteristic polynomial of a random unitary matrix and the Riemann zeta function*, PhD thesis (University of Bristol, 2001)

RAYMOND AND BEVERLY SACKLER SCHOOL OF MATHEMATICAL SCIENCES, TEL AVIV UNIVERSITY, TEL AVIV 69978, ISRAEL. CURRENT ADDRESS: AMERICAN INSTITUTE OF MATHEMATICS, 360 PORTAGE AVE., PALO ALTO, CA 94306-2244 ([hughes@aimath.org](mailto:hughes@aimath.org))